GUJARAT TECHNOLOGICAL UNIVERSITY

BE- SEMESTER 1st / 2nd EXAMINATION (OLD SYLLABUS) - SUMMER - 2017

Subject Code: 110009 Date:29/05/2017

Subject Name: Maths-II

Time: 2:30 PM to 05:30 PM Total Marks: 70

Instructions:

1. Attempt any five questions.

2. Make suitable assumptions wherever necessary.

3. Figures to the right indicate full marks.

Q.1 (a) (i) What conditions must b_1, b_2, b_3 satisfy in order for

 $x_1 + 2x_2 + 3x_3 = b_1, 2x_1 + 5x_2 + 3x_3 = b_2, x_1 + 8x_3 = b_3$ be consistent?

(ii) Obtain the reduced row echelon form of the matrix $A = \begin{bmatrix} 1 & 2 & 4 & 0 \\ -3 & 1 & 5 & 2 \\ -2 & 3 & 9 & 2 \end{bmatrix}$ 03

(b) Solve using Gauss-Elimination method 07

x + y + 2z = 8, -x - 2y + 3z = 1, 3x - 7y + 4z = 10

10 11 12 13 14 | 15 16 17 18 19 |

(ii) Is $T: R^3 \to R^3$ defined by T(x, y, z) = (x + 3y, y, z + 2x) linear? Is it one-to-one, onto or both? Justify.

(b) By Gauss-Jordan method find inverse of $\begin{bmatrix} 7 & 6 & 2 \\ -1 & 2 & 4 \end{bmatrix}$ 07

Q.3 (a) (i) Find a basis for the orthogonal complement of the subset of R^3 spanned by the vectors $v_1 = (1, -1, 3), v_2 = (5, -4, -4)$ and $v_3 = (7, -6, 2)$.

(ii) Let $W = \operatorname{Span}\left\{\left(\frac{4}{5}, 0, -\frac{4}{5}\right), (0, 1, 0)\right\}$. Express w = (1, 2, 3) in the form of **03**

 $w = w_1 + w_2$ where $w_1 \in W$ and $w_2 \in W^{\perp}$.

(b) Let R^3 have the Euclidean inner product. Use the Gram-Schmidt process to transform the basis vectors $u_1 = (1,1,1), u_2 = (-1,1,0)$ and $u_3 = (1,2,1)$ into orthonormal basis $\{v_1, v_2, v_3\}$ and show that the set $V = \{(1,0,0), (0,1,0), (0,0,1)\}$ is an orthonormal set in R^3 with Euclidean inner product.

- Q.4 (a)
- (i) Express the matrix $A = \begin{bmatrix} 2i & 3i & 4i \\ 0 & 2 & 5i \\ 2+i & 1-i & 0 \end{bmatrix}$ as a sum of Hermitian and Skew-

Hermitian matrices.

- (ii) Verify that the matrix A is unitary, where $A = \frac{1}{3}\begin{bmatrix} 2+i & 2i \\ 2i & 1-i \end{bmatrix}$
- **(b)** Consider the basis $S = \{v_1, v_2\}$ for R^2 , where $v_1 = (-2,1)$ and $v_2 = (1,3)$ and let $T: R^2 \to R^3$ be linear transformation such that $T(v_1) = (-1,2,0)$ and $T(v_2) = (0,-3,5)$. Find a formula for $T(x_1, x_2)$ and use that formula to find T(2,-3).
- Q.5 (a) (i) Show that the set of vectors 04 $\{2x^3 + x^2 + x + 1, x^3 + 3x^2 + x 2, x^3 + 2x^2 x + 3\}$ in P_3 is linearly independent.
 - (ii) Find basis and dimension of

$$W = \{ (a_1, a_2, a_3, a_4) \in R^4 / a_1 + a_2 = 0, a_2 + a_3 = 0, a_3 + a_4 = 0 \}$$

- **(b)** Check whether $V = R^2$ is a vector space with respect to the operations $(u_1, u_2) + (v_1, v_2) = (u_1 + v_1 2, u_2 + v_2 3)$ and $\alpha(u_1, u_2) = (\alpha u_1 + 2\alpha 2, \alpha u_2 3\alpha + 3), \alpha \in R$.
- Q.6 (a) (i) Determine which of the following is subspace or not (1) $V = \{(x, y) | x = 3y\}$ in R^2 (2) $V = \{(x, y) | x^2 = y^2\}$ in R^2
 - (ii) Determine whether the linear transformation is one-to-one or not $T: R^2 \to R^2$ where T(x, y) = (x + y, x y)
 - **(b)** Let $T_1: R^2 \to R^3, T_2: R^3 \to R^3, T_3: R^3 \to R^2$ be the linear transformation given by $T_1(x, y) = (-2y, 3x, x 2y), T_2(x, y, z) = (y, z, x), T_3(x, y, z) = (x + z, y z)$ find the domain and codomain of $(T_1 \circ T_2 \circ T_3)(x, y)$. Also find $(T_3 \circ T_2 \circ T_1)(1, 1)$.
- Q.7 (a)
 (i) Find eigen values and eigen vectors of the matrix $A = \begin{bmatrix} 3 & 1 & 4 \\ 0 & 2 & 6 \\ 0 & 0 & 5 \end{bmatrix}$.
 - (ii) Verify Cayley-Hamilton theorem for $A = \begin{bmatrix} 3 & 2 \\ 4 & -1 \end{bmatrix}$
 - **(b)** Find a matrix P that diagonalizes the matrix $A = \begin{bmatrix} 1 & 0 \\ -1 & 2 \end{bmatrix}$ and hence find A^{13} .

......