

Seat No.: \_\_\_\_\_

Enrolment No. \_\_\_\_\_

**GUJARAT TECHNOLOGICAL UNIVERSITY**  
**Diploma Engineering Semester –I/II Examination Jan. 2012**

**Subject code: 310034****Date: 16/01/2012****Subject Name: Mathematics-I****Time: 10.30 am – 01.00 pm****Total Marks: 70****Instructions:**

1. Attempt all questions.
2. Make suitable assumptions wherever necessary.
3. Figures to the right indicate full marks.
4. English version is considered Authentic.

**Q.1 (a)** Fill in the blanks (Not necessary to show the calculation) **10**

- (1)  $\log 1 \cdot \log 2 \cdot \log 3 \cdot \log 4 = \dots$ .
- (2)  $2^{-\log_2 3} = \dots$ .
- (3) If for G.P.,  $T_n = 3^n$  then common ratio  $r = \dots$ .
- (4) The Geometric Mean of 2 and 18 =  $\dots$ .
- (5) The 7<sup>th</sup> term in expansion of  $\left(x + \frac{1}{x}\right)^6 = \dots$ .
- (6) If  ${}^n C_5 = {}^n C_4$  then  $n = \dots$ .
- (7) If  $A = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ , then  $A^2 = \dots$ .
- (8) The value of  $\begin{vmatrix} \sec \theta & \tan \theta \\ \tan \theta & \sec \theta \end{vmatrix} = \dots$ .
- (9) If  $\bar{A} = \bar{i} - \bar{j} + \bar{k}$  then  $|\bar{A}| = \dots$ .
- (10) If  $\bar{a} = \bar{i} + \bar{j}$ ,  $\bar{b} = \bar{j} - \bar{k}$  then  $\bar{a} \cdot \bar{b} = \dots$ .

**(b)** (1) If the diameter of circle is 100cm and length of an arc is 25 cm, find the angle subtended by the arc at the centre of the circle. **02**

(2) Prove that,  $\cos \frac{\pi}{17} + \cos \frac{5\pi}{17} + \cos \frac{12\pi}{17} + \cos \frac{16\pi}{17} = 0$  **02**

**Q(2) (a)** Fill in the blanks (Not necessary to show the calculation) **07**

- (1)  $\cos \frac{\pi}{6} \cos \frac{\pi}{4} \cos \frac{\pi}{3} \cos \frac{\pi}{2} = \dots$ .
- (2)  $\sin^2 42^\circ + \sin^2 48^\circ = \dots$ .
- (3) The period of  $\tan 3x = \dots$ .
- (4) If  $\sin 15^\circ = \frac{\sqrt{6} - \sqrt{2}}{4}$  then  $\sin 165^\circ = \dots$ .
- (5) If  $\tan \theta = \frac{1}{3}$  then  $\sin 2\theta = \dots$ .
- (6) In usual notations for  $\Delta ABC$ , if  $a = 1$ ,  $b = 3$ ,  $c = 6$ , then  $s = \dots$
- (7) In any  $\Delta ABC$ , by second law of cosine(projection formula),

- (b) (1) If  $a^x = b^y = c^z$ , then prove that  $\log_a bc = x\left(\frac{1}{y} + \frac{1}{z}\right)$ , where **04**  
 $x, y, z \neq 0$ .
- (2) For a Geometrical Progression,  $T_5 = 81$  and  $T_2 = 3$ , find  $T_3$  and **03**  
 $S_3$ .

**OR**

- (b) (1) Solve the equation for x,  $\log_2(x+5) + \log_2(x-2) = 3$  **04**
- (2) Insert 5 Geometric Means between 320 and 5. **03**

- Q.3** (a) (1) Find middle term in the expansion of  $\left(\frac{x}{2} - \frac{2}{y}\right)^8$  **03**
- (2) Find the approximate value of  $\sqrt{102}$ , using binomial theorem. **02**
- (3) If  $A = \begin{bmatrix} 1 & 4 \\ 3 & 2 \\ 2 & 5 \end{bmatrix}$  and  $B = \begin{bmatrix} -1 & -2 \\ 0 & 5 \\ 3 & 1 \end{bmatrix}$ , Find  $3A - 2B$ . **02**
- (b) (1) Draw the graph of  $y = \cos \frac{x}{2}$ ,  $0 \leq x \leq 2\pi$ . **04**
- (2) Prove that :  $\tan 70^\circ - \tan 50^\circ - \tan 20^\circ = \tan 70^\circ \tan 50^\circ \tan 20^\circ$  **03**

**OR**

- Q.3** (a) (1) Find constant term in expansion of  $\left(x^2 - \frac{2}{x^2}\right)^8$ . **03**
- (2) Using binomial theorem expand  $(1-x)^{-\frac{1}{2}}$  up to first three terms. **02**
- (3) If  $A = \begin{bmatrix} 5 & -3 \\ 2 & 1 \end{bmatrix}$  then find  $A^{-1}$ . **02**
- (b) (1) Draw the graph of  $y = \sin x$ ,  $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$  **04**
- (2) Prove that :  $\tan 10^\circ + \tan 35^\circ + \tan 10^\circ \tan 35^\circ = 1$  **03**

- Q. 4** (a) (1) If  $A = \begin{bmatrix} 1 & 2 & 3 \\ 3 & -2 & 1 \\ 4 & 2 & 1 \end{bmatrix}$  find  $A^{-1}$  **03**
- (2) Show that the angle between the vectors  $i+2j$  and  $i+j+3k$  is **04**  
 $\sin^{-1} \sqrt{\frac{46}{55}}$
- (b) (1) Solve the following equations, by matrix method,  
 $3x + 2y = 7$  and  $11x - 4y = 3$ . **03**

(2) If  $\bar{a} = 3\bar{i} - \bar{j} - 4\bar{k}$ ,  $\bar{b} = -2\bar{i} + 4\bar{j} - 3\bar{k}$  and  $\bar{c} = \bar{i} + 2\bar{j} - \bar{k}$  then find the direction cosines of the vector  $3\bar{a} - 2\bar{b} + 4\bar{c}$ . 04

**OR**

**Q. 4** (a) (1) If  $A = \begin{bmatrix} -1 & 3 & 5 \\ 1 & -3 & -5 \\ -1 & 3 & 5 \end{bmatrix}$  then show that  $A^2 = A$ . 03

(2) Forces  $3\bar{i} - \bar{j} + 2\bar{k}$  and  $\bar{i} + 3\bar{j} - \bar{k}$  act on a particle and particle moves from  $2\bar{i} + 3\bar{j} + \bar{k}$  to  $5\bar{i} + 2\bar{j} + \bar{k}$  under these forces. Find the work done by the forces. 04

(b) (1) If  $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$ , then prove that  $A^2 - 5A - 2I = 0$ . 03

(2) Find a unit vector perpendicular to the plane of vector  $\bar{a} = 5\bar{i} + 7\bar{j} - 2\bar{k}$  and  $\bar{b} = 3\bar{i} + \bar{j} - 2\bar{k}$ . 04

**Q. 5** (a) (1) Prove that  $\frac{\cos A + \cos 3A + \cos 5A}{\sin A + \sin 3A + \sin 5A} = \cot 3A$  04

(2) Prove that  $\tan^{-1}\left(\frac{1}{2}\right) + \tan^{-1}\left(\frac{1}{3}\right) = \frac{\pi}{4}$  03

(b) (1) In usual notations for  $\Delta ABC$  prove that  $a = b \cos C + c \cos B$  04

(2) For  $\Delta ABC$ ,  $a = 7$ ,  $b = 4\sqrt{3}$  and  $c = \sqrt{13}$ , find the measure of the smallest angle of the triangle. 03

**OR**

**Q. 5** (a) (1) Prove that,  $\frac{1 + \sin 2\theta - \cos 2\theta}{1 + \sin 2\theta + \cos 2\theta} = \tan \theta$  04

(2) Prove that  $2 \tan^{-1}\left(\frac{1}{3}\right) + \tan^{-1}\left(\frac{1}{7}\right) = \frac{\pi}{4}$  03

(b) (1) In usual notations for  $\Delta ABC$ , prove that  $b = a \cos C + c \cos A$ . 04

(2) Solve the  $\Delta ABC$ , having  $A = 30^\circ$ ,  $b = 1$  and  $c = \sqrt{3}$ . 03

\*\*\*\*\*

પ્ર 1 (અ) ખાલી જગ્યા પૂરો .(ગણતરી દર્શાવવી જરૂરી નથી )

10

(1)  $\log 1 \cdot \log 2 \cdot \log 3 \cdot \log 4 = \dots$

(2)  $2^{-\log_2 3} = \dots$

(3) એક સમગુણોતર શ્રેણી માટે ,જો  $T_n = 3^n$ , તો સામાન્ય ગુણોતર  $r = \dots$ .

(4) 2 અને 18 નો સમગુણોતર મધ્યક =-----

(5)  $\left( x + \frac{1}{x} \right)^6$  ના વિસ્તરણ માં સાતમું પદ =-----.

(6) જો  ${}^n C_5 = {}^n C_4$  તો  $n = \dots$

(7) જો  $A = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ , તો  $A^2 = \dots$ .

(8)  $\begin{vmatrix} \sec \theta & \tan \theta \\ \tan \theta & \sec \theta \end{vmatrix} = \dots$ .

(9) જો  $\bar{A} = \bar{i} - \bar{j} + \bar{k}$  તો  $|\bar{A}| = \dots$ .

(10) જો  $\bar{a} = \bar{i} + \bar{j}$ ,  $\bar{b} = \bar{j} - \bar{k}$  તો  $\bar{a} \cdot \bar{b} = \dots$ .

(બ) (1) જો એક વર્તુળ ના વ્યાસ ની લંબાઈ 100cm અને તેના એક ચાપ ની લંબાઈ 25 cm, હોય તો ચાપે કેન્દ્ર આગળ આંતરેલો ખુણો શોધો.

2) સાબિત કરો કે ,  $\cos \frac{\pi}{17} + \cos \frac{5\pi}{17} + \cos \frac{12\pi}{17} + \cos \frac{16\pi}{17} = 0$  02

પ્ર.2 (અ) ખાલી જગ્યા પૂરો .(ગણતરી દર્શાવવી જરૂરી નથી ) 07

(1)  $\cos \frac{\pi}{6} \cos \frac{\pi}{4} \cos \frac{\pi}{3} \cos \frac{\pi}{2} = \dots$ .

(2)  $\sin^2 42^\circ + \sin^2 48^\circ = \dots$ .

(3)  $\tan 3x$  નો આવર્તમાન =-----.

(4) જો  $\sin 15^\circ = \frac{\sqrt{6} - \sqrt{2}}{4}$ , હોય તો  $\sin 165^\circ = \dots$ .

(5) જો  $\tan \theta = \frac{1}{3}$  તો  $\sin 2\theta = \dots$ .

(6) પ્રયત્નિત સંકેત પ્રમાણે ,જો  $\Delta ABC$ , માં  $a = 1$ ,  $b = 3$ ,  $c = 6$  , હોય તો  $s = \dots$

(7) કોઇ પણ  $\Delta ABC$  માં કોસાઇન ના બીજા નિયમ (પ્રક્ષેપ સુત્ર )  
પ્રમાણે  $c = \dots$

(બ) (1) જો  $a^x = b^y = c^z$ , હોય તો સાબિત કરો કે 04

$$\log_a bc = x \left( \frac{1}{y} + \frac{1}{z} \right),$$
 જ્યાં  $x, y, z \neq 0.$

(2) કોઇ પણ સમગુણોતર શ્રેણી માં ,  $T_5 = 81$  અને  $T_2 = 3$ , હોય તો  $T_3$  અને  $S_3$ .શોધો 03

- (બ) (1) ની કિમંત માટે સમીકરણ ઉકેલો,  $\log_2(x+5) + \log_2(x-2) = 3$  04  
 (2) 320 અને 5 વચ્ચે 5 સમગુણોતર મધ્યકો મૂકો . 03

- પ્ર 3 (અ) (1)  $\left(\frac{x}{2} - \frac{2}{y}\right)^8$  ના વિસ્તરણ માં મધ્યમ પદ શોધો. 03  
 (2) દ્વિપદી પ્રમેય નો ઉપયોગ કરી ને  $\sqrt{102}$  , ની આશરે કીમત શોધો 02  
 (3) જો  $A = \begin{bmatrix} 1 & 4 \\ 3 & 2 \\ 2 & 5 \end{bmatrix}$  અને  $B = \begin{bmatrix} -1 & -2 \\ 0 & 5 \\ 3 & 1 \end{bmatrix}$ , હોય તો  $3A - 2B$  શોધો 02  
 (બ) (1)  $y = \cos \frac{x}{2}$  નો આલેખ દોરો , (  $0 \leq x \leq 2\pi$  ) 04  
 (2) સાબિત કરો કે:  $\tan 70^\circ - \tan 50^\circ - \tan 20^\circ = \tan 70^\circ \tan 50^\circ \tan 20^\circ$  03

અથવા

- પ્ર.3 (અ) (1)  $\left(x^2 - \frac{2}{x^2}\right)^8$  , ના વિસ્તરણ માં અચલ પદ શોધો 03  
 (2)  $(1-x)^{-\frac{1}{2}}$  નો દ્વિપદી પ્રમેય થી પ્રથમ ત્રણ પદો સુધી વિસ્તરણ કરો 02  
 (3) જો  $A = \begin{bmatrix} 5 & -3 \\ 2 & 1 \end{bmatrix}$  તો  $A^{-1}$  શોધો . 02  
 (બ) (1)  $y = \sin x$ , નો આલેખ દોરો , (  $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$  ) 04  
 (2) સાબિત કરો કે :  $\tan 10^\circ + \tan 35^\circ + \tan 10^\circ \tan 35^\circ = 1$  03

- પ્ર 4 (અ) (1) જો  $A = \begin{bmatrix} 1 & 2 & 3 \\ 3 & -2 & 1 \\ 4 & 2 & 1 \end{bmatrix}$  તો  $A^{-1}$  શોધો 03  
 (2) સાબિત કરો કે . સદિશો  $i+2j$  , અને  $i+j+3k$  વચ્ચે નો ખૂણો 04  
 $\sin^{-1} \sqrt{\frac{46}{55}}$  છે
- (બ) (1) શ્રેણીક ની મદદ થી નીચે ના સમીકરણો નો ઉકેલ મેળવો 03  
 $3x + 2y = 7$  અને  $11x - 4y = 3$ .

- (2) જો  $\bar{a} = 3\bar{i} - \bar{j} - 4\bar{k}$ ,  $\bar{b} = -2\bar{i} + 4\bar{j} - 3\bar{k}$  અને  $\bar{c} = \bar{i} + 2\bar{j} - \bar{k}$  04,  
 હોય તો સદિશ  $3\bar{a} - 2\bar{b} + 4\bar{c}$ . ના દિક્કોસાઇન મેળવો

અથવા

પ્ર 4 (અ) જો  $A = \begin{bmatrix} -1 & 3 & 5 \\ 1 & -3 & -5 \\ -1 & 3 & 5 \end{bmatrix}$  હોય તો સાબિત કરો કે  $A^2 = A$ .

(2) એક કણ પર લાગતા બળો  $3\bar{i} - \bar{j} + 2\bar{k}$  અને  $\bar{i} + 3\bar{j} - \bar{k}$ ,  
તે કણ ને  $2\bar{i} + 3\bar{j} + \bar{k}$  થી  $5\bar{i} + 2\bar{j} + \bar{k}$  સુધી ખસેડે છે તો તે  
બળો થી થયેલું કાર્ય શોધો

(અ) (1) જો  $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$ , હોય તો સાબિત કરો કે  $A^2 - 5A - 2I = 0$ .

(2) સદિશ  $\bar{a} = 5\bar{i} + 7\bar{j} - 2\bar{k}$ , અને  $\bar{b} = 3\bar{i} + \bar{j} - 2\bar{k}$ . પર લંબ એકમ  
. સદિશ મેળવો

પ્ર 5 (અ) (1) સાબિત કરો કે  $\frac{\cos A + \cos 3A + \cos 5A}{\sin A + \sin 3A + \sin 5A} = \cot 3A$

(2) સાબિત કરો કે  $\tan^{-1}\left(\frac{1}{2}\right) + \tan^{-1}\left(\frac{1}{3}\right) = \frac{\pi}{4}$

(અ) (1) પ્રચલિત સંકેતો માં  $\Delta ABC$  માટે સાબિત કરો કે,  
 $a = b \cos C + c \cos B$

(2)  $\Delta ABC$  માટે  $a = 7$ ,  $b = 4\sqrt{3}$  અને  $c = \sqrt{13}$ , હોય તો સૌથી નાના  
ખુણા નું માપ શોધો.

## અથવા

(અ)

પ્ર 5 (1) સાબિત કરો કે,  $\frac{1 + \sin 2\theta - \cos 2\theta}{1 + \sin 2\theta + \cos 2\theta} = \tan \theta$

(2) સાબિત કરો કે,  $2 \tan^{-1}\left(\frac{1}{3}\right) + \tan^{-1}\left(\frac{1}{7}\right) = \frac{\pi}{4}$

(અ)

(1) પ્રચલિત સંકેતો માં  $\Delta ABC$ , માટે સાબિત કરો કે  
 $b = a \cos C + c \cos A$ .

(2) જો  $\Delta ABC$  માં  $A = 30^\circ$ ,  $b = 1$  અને  $c = \sqrt{3}$  હોય તો  
 $\Delta ABC$  નો ઉકેલ શોધો

\*\*\*\*\*